

References

de Costa, SP., van der Schans, CP., Boelema, SR., van der Meij E., Boeman, M, Bos, AF. (2010). Sucking Patterns in full-term infants between birth and 10 weeks of age. *Infant Behavior and Development*. 33(1): 61-67.

Freyer, FJ. www.bostonglobe.com “Hospitals adapt as opioid epidemic hits infant victims.” published June 28,2015.

Hall, et al. (2016). Clinical management of breast feeding Mother- Infant Dyad in recovery from opioid dependency. *Journal of Addictions Nursing*. 27(2), page(s) 68-77.

Jansson, LM., Valez, M., Harrow, C. (2009) The Opioid Exposed Newborn: Assessment and Pharmacological Management.Journal of Opioid Management.5(1):47-55.

Kocherlakota, Prabhakar.(2014). Neonatal Abstinence Syndrome. *Pediatrics*. 134(2), pages 547-561.

Lester, BM., Tronick, EZ. (2004). The Neonatal Intensive Care Unit Network Neurobehavioral Scale Procedures. *Pediatrics*. 113(3),pages 641- 667.

Pados, BF., Park, J., Thoyre, SM, Estrem, H., Nix, WB. (2015). Milk Flow Rates From Bottle Nipple Used for Feeding Infants Who Are Hospitalized. *American Journal of Speech Language Pathology*, 1;24(4): 671-679.

Patrick, SW., Schumacher, RE., Horbar, JD., Buss-Frank, ME., Edwards, EM., Morrow, KA., Ferrelli, KR., Pincarillo, AP., Gupta, M., Soil, RF. (2016). Improving Care for Neonatal Abstinence Syndrome. *Pediatrics*. e20153835.

Patrick SW, et al. (2012).Neonatal Abstinence Syndrome and Associates Healthcare Expenditures – United States, 2000-2009. *JAMA*.May 9;307(18):1934-40.

Ross, ES,Philibin, MK. (2011). Supporting oral feeding in fragile infants: an evidence- based method for quality bottle feedings of preterm, ill, and fragile infants. *Journal of Perinatal and Neonatal Nursing*, 25(4): 349-57.

Ross, ES. SOFFI Training Course, National Children's Hospital, April 2016
(powerpoint slide)

Thoyre, SM., Shaker,CS., Pridham,KF. The Early Feeding Skills Assessment for Preterm Infants. *Neonatal Network*. 24(3); 7-16.